

Controlling medusahead with high rates of *Milestone*

G.B. Kyser
V.F. Peterson
J.S. Davy
J.M. DiTomaso

Medusahead

(*Taeniatherum caput-medusae*)



Medusahead

(*Taeniatherum caput-medusae*)




Medusahead

(*Taeniatherum caput-medusae*)



Medusahead


(*Taeniatherum caput-medusae*)



Control seed
production

Burn it, eat it,
spray it with *Plateau*





The background of the image shows a field of yellow thistle flowers (Carduus pumilus) growing in a dry, open landscape. In the distance, there are rolling hills covered with sparse vegetation. The foreground is dominated by the spiny, yellow-flowered thistle plants.


Milestone®

(aminopyralid, 2 lb ae/gal)

- Broadleaf selective (auxinic)
- Mostly used for thistles
- Pre- and postemergence
- 3 to 7 oz/acre
(14 oz for spot treatment)












Postemergence –
14 oz *Milestone*






Preemergence –
3.5 oz *Milestone*




Preemergence –
7 oz Milestone



Preemergence –
14 oz Milestone







Google earth

Eye alt 323.96 km

Preemergence treatments

late September to early October 2009

Herbicide	Active ingredient	Rates (oz / acre)
Milestone	aminopyralid (2 lb/gal)	0, 3, 5, 7, 14
Matrix	rimsulfuron (25%)	1, 2
Plateau	imazapic (2 lb/gal)	8

Plots 10 ft by 30 ft


Treatments in four replications X three sites

CO₂ backpack sprayer, 10-ft boom with six 8002 nozzles, 20 gpa

Evaluations

Data collected	Sampling units per plot	Timing
Vegetative cover	Three 1-m ² quadrats	Late May 2010
Biomass	Three 0.25-m ² quadrats	Late May 2010
Medusahead seedheads for viability	10 heads → 50 seeds	Mid-June 2010




















2011 to 2012 – Red Bluff

14 oz *Milestone* in October – 80% control

2011 to 2012 – Red Bluff

14 oz *Milestone* in October – 80% control

Split application (7 oz Oct, 7 oz Jan)

2011 to 2012 – Red Bluff

14 oz *Milestone* in October – 80% control

Split application (7 oz Oct, 7 oz Jan) – 97% control

2011 to 2012 – Red Bluff

14 oz *Milestone* in October – 80% control

Split application (7 oz Oct, 7 oz Jan) – 97% control

Maximum Application Rate: On all labeled use sites do not broadcast apply more than 7 fl oz per acre of *Milestone* per year. ... Spot treatments may be applied at an equivalent broadcast rate of up to 14 fl oz per acre per annual growing season; however, not more than 50% of an acre may be treated at that rate.

Large-scale test

14 oz/acre

Milestone

near Red Bluff



Acknowledgements

Winters: thanks to Morgan Doran (CE Farm Advisor, Solano, Yolo, and Napa counties), to the Audubon Bobcat Ranch, and to the Lindemann family for allowing access to the site.

Marysville: thanks to the Sierra Foothill Research and Extension Center and to Dustin Flavell (Center Superintendent).

Red Bluff: thanks to the Gallatin Ranch.

And thanks to Carlos Figueroa and Joe Webster, undergraduate research assistants, for processing the samples and performing the germination tests.

LITERATURE CITED

- Bovey, R. W., D. LeTourneau, and L. C. Erickson. 1961. The chemical composition of medusahead and downy brome. *Weeds* 9:307-311.
- Carrithers, V. F., P. L. Burch, W. N. Kline, R. A. Masters, J. A. Nelson, M. B. Halstvedt, J. L. Troth, and J. M. Breuninger. 2005. Aminopyralid: a new active ingredient for control of broadleaf invasive and noxious weeds. Abstracts: Weed Science Society of America, 45th Annual Meeting, 7-10 February 2005; Honolulu, HI, USA: p. 114.
- DiTomaso, J. M., G. B. Kyser, M. R. George, M. P. Doran, and E. A. Laca. 2008. Control of medusahead using timely sheep grazing. *Invasive Plant Science and Management* 1:241-247.
- DiTomaso, J. M., G. B. Kyser, and M. S. Hastings. 1999a. Prescribed burning for control of yellow starthistle (*Centaurea solstitialis*) and enhanced native plant diversity. *Weed Science* 47:233-242.
- DiTomaso, J. M., G. B. Kyser, S. B. Orloff, S. F. Enloe, and G. A. Nader. 1999b. New growth regulator herbicide provides excellent control of yellow starthistle. *California Agriculture* 53:12-16.
- Duncan, C. L., and J. K. Clark. 2005. Invasive plants of range and wildlands and their environmental, economic, and societal impacts. Lawrence, KS, USA: Weed Science Society of America. 222 p.
- Enloe, S. F., G. B. Kyser, S. A. Dewey, V. Peterson, J. M. DiTomaso. 2008. Russian knapweed (*Acroptilon repens*) control with low rates of aminopyralid on range and pasture. *Invasive Plant Science and Management* 1:385-389.
- Enloe, S. F., R. G. Lym, R. Wilson, P. Westra, S. Nissen, G. Beck, M. Moechnig, V. Peterson, R. A. Masters, and M. Halstvedt. 2007. Canada thistle (*Cirsium arvense*) control with aminopyralid in range, pasture, and noncrop areas. *Weed Technology* 21:890-894.
- Ferrell, J. A., J. J. Mullahey, K. A. Langeland, and W. N. Kline. 2006. Control of tropical soda apple (*Solanum viarum*) with aminopyralid. *Weed Technology* 20:453-457.
- George, M. R. 1994. Annual rangeland management principles and practices. In S. B. Monsen and S. G. Kitchen [COMPS.]. Proceedings: Symposium on ecology, management and restoration of intermountain annual rangelands, 18-21 May 1992; Boise, ID, USA. Gen. Tech. Rep. INT-313. US Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, UT, USA. p. 392-395.
- Harris, C. A. 1977. Root phenology as a factor of competition among grass seedlings. *Journal of Range Management* 30:172-177.
- Hironaka, M. 1961. The relative rate of root development of cheatgrass and medusahead. *Journal of Range Management* 14:263-267.
- Hulting, A. G., J. L. Haavisto, and G. Pirelli. 2012. Biology and management of hare barley in cool-season grass pastures. Proceedings of the 65th Western Society of Weed Science Conference; 12-15 March 2012; Reno, NV, USA (in press).
- Kyser, G. B., J. M. DiTomaso, M. P. Doran, S. B. Orloff, R. G. Wilson, D. L. Lancaster, D. F. Lile, and M. L. Porath. 2007. Control of medusahead (*Taeniatherum caput-medusae*) and other annual grasses with imazapic. *Weed Technology* 21:65-75.
- Kyser, G. B., M. P. Doran, N. K. McDougald, S. B. Orloff, R. N. Vargas, R. G. Wilson, and J. M. DiTomaso. 2008. Site characteristics determine the success of prescribed burning for medusahead control. *Invasive Plant Science and Management* 1:376-384.
- Monaco, T. A., T. M. Osmond, and S. A. Dewey. 2005. Medusahead control with fall- and spring-applied herbicides on northern Utah foothills. *Weed Technology* 19:653-658.
- Rinella, M. J., M. R. Haferkamp, R. A. Masters, J. M. Muscha, S. E. Bellows, and L. T. Vermeire. 2010. Growth regulator herbicides prevent invasive annual grass seed production. *Invasive Plant Science and Management* 3:12-16.
- SAS Institute. 2008. JMP statistics and graphics guide. Version 8.0. Cary, NC, USA: SAS Institute, Inc. 1230 p.
- Seefeldt, S. and J. Conn. 2011. Control of orange hawkweed (*Hieracium aurantiacum*) in southern Alaska. *Invasive Plant Science and Management* 4:87-94.
- Sheley, R. L., M. F. Carpinelli, and K. J. Reever Morgan. 2007. Effects of imazapic on target and nontarget vegetation during revegetation. *Weed Technology* 21:1071-1081.
- Shinn, S. L. and D. C. Thill. 2002. The response of yellow starthistle (*Centaurea solstitialis*), annual grasses, and smooth brome (*Bromus inermis*) to imazapic and picloram¹. *Weed Technology* 16:366-370.
- Sweet, S. B., G. B. Kyser, and J. M. DiTomaso. 2008. Susceptibility of exotic annual grass seeds to fire. *Invasive Plant Science and Management* 1:158-167.
- UC IPM. 2010. UC IPM online. Available at <http://www.ipm.ucdavis.edu/WEATHER/index.html>. Accessed 1 December 2010.
- USDA, NRCS. 2011. Web soil survey. Available at <http://websoilsurvey.nrcs.usda.gov/app/>. Accessed 6 July 2011.
- Young, J. A., R. A. Evans, and J. Robison. 1972. Influence of repeated annual burning on a medusahead community. *Journal of Range Management* 25:372-375.