

## Field Guide for Managing Cheatgrass in the Southwest










Repeated mowing every 2 to 3 weeks during spring and summer may be as effective as an application of glyphosate.

However, mowing just a single time has not been found to be effective in preventing seed production since plants often rapidly regrow and still produce seed.











## syngenta

## Herbicide

For the control of grass weeds in andscape areas, roadsides, nurseries, greenhouses, flower beds, groundcovers, interiorscapes, parks, sports fields, golf courses, commercial and residential areas.

Active Ingredient:

Fluazifop-P-butyl

Butyl (R)-2-[4-[[5-(trifluoromethyl)-2-

pyridinyl]oxy]phenoxy]propanoate\* . . . . . . . 24.5%

Other Ingredients 75.5%

*Total:* 100.0%

\*Fusilade II Turf and Ornamental Herbicide is an emulsifiable concentrate containing 2 pounds (+) isomer (fluazifop-P-butyl) per gallon.

Contains petroleum distillate.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-1084 EPA Est. No. 11773-IA-01<sup>VWC</sup>

EPA Est. No. 46073-TN-003<sup>NTM</sup>

TARGET PESTS: Invasive annual grasses – Barbed Goatgrass – Aegilops and Cheatgrass – Bromus tectorum

Material (Add to tank in order)	Rate per 100 gal	Rate per gal (handhelds)	Volume/Acre
(1) Fusilade II Herbicide – 100-1084	0.5 gallon	0.67 oz.	Variable – Low Volume Spot Spray Treatments
(2a) Renegade-EA Surfactant – 2935-50201	1 gallon	1.3 oz.	
Or (2b) Competitor ESO Surfactant- 2935-50173	1 gallon	1.3 oz.	
(3) Hi-Light Blue Dye (if necessary)	1 quart	0.3 oz.	

